

DOI: 10.3969/j.issn.1000-3142.2014.03.001

王文采. 东川铁线莲, 云南毛茛科一新种[J]. 广西植物 2014, 34(3): 287–289

WANG Wen-Tsai. *Clematis dongchuanensis*, a new species of Ranunculaceae from Yunnan [J]. *Guizhou Botany* 2014, 34(3): 287–289

## 东川铁线莲, 云南毛茛科一新种

王文采

(中国科学院植物研究所 系统与进化植物学国家重点实验室, 北京 100093)

**摘要:** 描述了自云南东北部发现的毛茛科铁线莲属一新种 东川铁线莲。此种由于其叶为单叶, 萼片直立, 雄蕊的花丝和花药均被短柔毛而与曲柄铁线莲近缘, 与后者的区别在于其叶呈心状卵形, 叶缘具粗牙齿, 花序为二歧聚伞花序, 以及花药顶端钝, 不具短尖头。

**关键词:** 毛茛科; 铁线莲属; 新种; 云南

中图分类号: Q949.746 文献标识码: A 文章编号: 1000-3142(2014)03-0287-03

## *Clematis dongchuanensis*, a new species of Ranunculaceae from Yunnan

WANG Wen-Tsai

(State Key Laboratory of Systematic and Evolutionary Botany, Institute of Botany,  
the Chinese Academy of Sciences, Beijing 100093, China)

**Abstract:** A species of the genus *Clematis* (Ranunculaceae) *C. dongchuanensis* is described as new from northeastern Yunnan. In having simple leaves, erect sepals and puberulous filaments and anthers of stamens this species is related to *C. repens* Finet & Gagnep., and from the latter differs in its cordate-ovate and coarsely dentate leaves, dichotomous cymes and obtuse, not apiculate apexes of anthers.

**Key words:** Ranunculaceae; *Clematis*; new species; Yunnan

### 东川铁线莲 图1

***Clematis dongchuanensis*** W. T. Wang, sp.

nov. Fig. 1.

sect. *Viorna* (Reichb.) Prantl subsect. *Connatae* Koehne

Ob folia simplicia, sepala erecta, et staminum filaments antherasque omnino puberulas species nova haec est affinis *C. repenti* Finet & Gagnep. quae foliis simplicibus vel folii ternati foliolis ellipticis indivisis basi late cuneatis vel rotundatis raro subcordatis margine sparse denticulatis vel integris, cymis axillaribus 1-flo-ris, sepalis dorso glabris, antheris apice apiculatis facile

differat.

Woody vine. Stem 1–2 m long (according to the collector); branchlets slender ca. 1 mm across, sparsely puberulous. Leaves opposite, simple; blades papery, cordate-ovate or pentagonal-ovate 9–12 cm long, 4–6.5 cm broad, apex acuminate, base cordate, 3-lobed; central lobe larger, ovate, 6–8 cm long, with margin unequally coarsely dentate, lateral lobes smaller, obliquely flabellate, unequally 2-lobulate, with margin unequally coarsely dentate; surfaces adaxially sparsely strigillose, abaxially appressed-pubescent; venation trinerved, with nerves nearly flat; petioles 4–6.5 cm long, sparsely puberulous.

收稿日期: 2013-03-20 修回日期: 2013-08-15

作者简介: 王文采(1926-), 男, 山东掖县人, 研究员, 中国科学院院士, 著名植物分类学家, 长期从事有花植物学研究。

Dichotomous cyme terminal (?) ca. 15 cm broad three times branched ca. 12-flowered;




图 1 东川铁线莲 A. 部分枝条; B. 聚散花序; C. 萼片(背面观); D. 雄蕊; E. 心皮。(孙英宝绘图)

Fig. 1 *Clematis dongchuanensis* A. Part of a branch; B. Cyme; C. Sepal (dorsal view); D. Stamen; E. Carpels. (Drawn by SUN Ying-Bao)

peduncle ca. 7 cm long subglabrous; bracts 2 opposite, lanceolate-linear ca. 8 mm long, 1 mm broad, abaxially sparsely pubescent, base decurrent and auriculate; branches and pedicels glabrous; bracts of branches opposite, linear 2.5–5.8 mm long puberulous; pedicels 0.35–6.5 cm long. Flower slightly pendulous: Calyx broadly campanulate; sepals 4, erect, yellow-green (according to the collector), oblong-ovate, 13–15 mm long, 5–6 mm broad, adaxially and abaxially densely appressed-puberulous, apex obtuse, margin above velutinous. Stamens ca. 50, ca. 11–13.2 mm long; filaments narrow-linear 9–10 mm long, ca. 0.4 mm broad, densely puberulous, longitudinally 1-nerved; anthers linear 2.2–2.8 mm long, 0.4 mm broad, abaxially puberulous, apex obtuse. Carpels ca. 40, ca. 9 mm long; ovaries ca. 1 mm long, appressed-pubescent; styles slender, ca. 8 mm long, below densely long sericeous (with hairs 2–4.5 mm long) above densely sericeous-pubescent (with hairs 0.5–1 mm long) apex slightly thickened.

木质藤本。茎长1–2 m (据采集者); 小枝细, 粗约1 mm, 疏被短柔毛。叶对生, 为单叶; 叶片纸质, 心状卵形或五角状卵形, 长9–12 cm, 宽4–6.5 cm, 顶端渐尖, 基部心形, 3浅裂, 中央裂片较长, 卵形, 长6–8 cm, 边缘有不等大的粗牙齿, 侧裂片较小, 斜扇形, 不等微2裂, 边缘有不等大的粗牙齿, 上面疏被短糙伏毛, 下面被贴伏柔毛, 三出脉, 脉近平; 叶柄长4–6.5 cm, 疏被短柔毛。二歧聚伞花序顶生(?)，宽约15 cm, 三回分枝, 约有12花; 花序梗长约7 cm, 近无毛; 苞片2, 对生, 披针状条形, 长约8 mm, 宽1 mm, 下面疏被柔毛, 基部下延, 呈耳状; 分枝和花梗无毛; 分枝的苞片对生, 条形, 长2.5–5.8

mm, 被短柔毛; 花梗长0.35–6.5 cm。花稍下垂: 花萼宽钟状; 萼片4, 直立, 黄绿色(据采集者), 长圆状卵形, 长13–15 mm, 宽5–6 mm, 两面密被贴伏短柔毛, 顶端钝, 边缘上部被短绒毛。雄蕊约50, 长11–13.2 mm; 花丝狭条形, 长9–10 mm, 宽约0.4 mm, 密被短柔毛, 具1条纵脉; 花药条形, 长2.2–2.8 mm, 宽0.4 mm, 背面被短柔毛, 顶端钝。心皮约40, 长约9 mm; 子房约1 mm, 被贴伏柔毛; 花柱细长, 长约8 mm, 下部密被长绢毛(毛长2–4.5 mm), 上部密被绢状柔毛(毛长0.5–1 mm), 顶端稍变粗。

云南(Yunnan): 东川(Dongchuan), 因民(Yinmin) 小山脑(Xiaoshannao) alt. 2290 m, 崖边(by cliff), 藤本长1–2 m (vine 1–2 m long), 花黄绿色(fls. yellow-green), 1984-09-11, 蓝顺彬(Lan Shun-bin) 217 (holotype PE).

本种的叶为单叶, 萼片直立, 雄蕊花丝和花药均被短柔毛, 曲柄铁线莲(*Clematis repens* Finet & Gagnep.)也具上述形态特征, 这种情况说明本种在亲缘关系上接近曲柄铁线莲, 与后者的区别是本种的叶呈心状卵形, 3浅裂, 边缘有粗牙齿, 花序为具约12朵花的二歧聚伞花序, 萼片两面密被短柔毛, 雄蕊花药顶端钝, 无短尖头。在曲柄铁线莲, 其单叶或有时为三出复叶的小叶均呈椭圆形, 不分裂, 基部宽楔形或圆形, 稀浅心形, 边缘具少数小牙齿或全缘, 聚伞花序腋生, 只具1花, 萼片外面无毛, 雄蕊花药顶端有短尖头(方明渊, 1980)。

致谢 深切感谢孙英宝先生为本文绘图。

#### 参考文献:

Fang MY(方明渊). 1980. *Clematis* sect. *Viorna* [M] // Flora of China(中国植物志). Beijing(北京): Science Press(科学出版社) 28: 88–133

(上接第332页 Continue from page 332)

munity structure in a forest ecosystem in the Southern Mountains of Urumqi (乌鲁木齐南部山区森林生态系统树生地衣群落) [J]. *Chin Bull Bot*(植物学报) 44(5): 578–586  
Tumur A(艾尼瓦尔·吐米尔), Abbas A(阿不都拉·阿巴斯). 2006. Floor lichen diversity under different vegetation types in Two-river sources Natural Reserve in Altay Mountains, Xinjiang (新疆阿勒泰山两河源自然保护区地面生地衣的物种多样性) [J]. *Biodivers Sci*(生物多样性) 14(5): 444–450  
Tumur A(艾尼瓦尔·吐米尔), Abdulla A(阿地力江·阿不都拉), Abbas A(阿不都拉·阿巴斯). 2011. Distribution pattern of floor lichen communities in Southern mountainous area of Urumqi, Xinjiang, China(乌鲁木齐南部山区地生地衣群落分布格

局) [J]. *Chin Biodivers*(生物多样性) 19(5): 574–580  
Wang YL(王玉良), Abdulla A(阿地力江·阿不都拉), Tumur A(艾尼瓦尔·吐米尔), et al. 2012. Diversity and community quantitative characteristics of floor lichen in Yaoluoping National Nature Reserve of Anhui, China(鹞落坪自然保护区地面生地衣多样性及群落数量特征) [J]. *Plant Divers Resour*(植物分类与资源学报) 34(2): 199–207  
Yang HX(阳含熙), Lu ZY(卢泽愚). 1981. Methodology of Quantitative Taxonomy in Plant Ecology(植物生态学数量分析方法) [M]. Beijing: Sciences Press: 90–252  
Zedda L, Rambold G. 2009. Diversity and ecology of soil lichens in the Knersvlakte(South Africa) [J]. *Bryologist* 112: 19–29