

DOI: 10.3969/j.issn.1000-3142.2014.01.001

王文采 柔毛赤车 广西荨麻科一新种[J]. 广西植物 2014,34(1):1-3

Wang WT. *Pellionia mollissima* a new species of Urticaceae from Guangxi [J]. Guihaia 2014,34(1):1-3

## 柔毛赤车，广西荨麻科一新种

王文采

(中国科学院植物研究所 系统与进化植物学国家重点实验室,北京 100093)

摘要：描述了自广西西南部发现的荨麻科(Urticaceae)赤车属(*Pellionia*)一新种柔毛赤车(*P. mollissima*)。

此种的特征是其瘦果具短线纹和瘤状突起,据此果实特征,可将此种与赤车属的其他种相区别。

关键词：荨麻科；赤车属；新种；广西

中图分类号：Q949 文献标识码：A 文章编号：1000-3142(2014)01-0001-03

## *Pellionia mollissima*, a new species of Urticaceae from Guangxi

WANG Wen-Tsai

(State Key Laboratory of Systematic and Evolutionary Botany, Chinese Academy of Science, Beijing 100093, China)

**Abstract:** A species of the genus *Pellionia* (Urticaceae) *P. mollissima* is described as new from southwestern Guangxi. This species is characterized by its lineolate and tuberculate achenes and by this unique fruit character it may be distinguished from all the other species of *Pellionia*.

**Key words:** Urticaceae; *Pellionia*; new species; Guangxi

### 柔毛赤车 图 1

#### *Pellionia mollissima* W. T. Wang sp. nov. Fig.

1.

Species nova haec achenis longitudinaliter lineolatis et minutissime tuberculatis insignis est, secundum fructus characterem hunc a speciebus omnibus ceteris generis *Pellioniae* diffeat, etiam foliis utrinque densissime puberulis a speciebus sinensis ceteris generis hujus recedit.

Perennial herbs, turns black when drying. Stems ca. 65 cm tall, near base 6 mm across, above densely puberulous (with hairs 0.2 mm long), below glabrous and longitudinally shallowly 10-sulcate. Leaves sessile; blades thin papery, obliquely oblandeolate or long elliptic, 13-17 cm long, 2-4.8 cm broad, apex long acuminate, base at leaf narrow side long attenuate and at broad side obtuse or auriculate, margin denticulate; surfaces adaxially and abaxially very densely puberulous (with hairs 0.2-0.5 mm long); venation semi-triplinerved, with lateral nerves 3 at leaf narrow side, 5 at broad side; cystoliths dense, bacilliform, 0.2-0.5 mm long; stipules membranous, narrow-lanceolate or linear, 9-11.5 mm long, 2-2.8 mm broad, abaxially puberulous. Staminate cymes unknown. Infructescences singly axillary, small, ca. 3 mm long and broad, sessile, below from base very shortly 4-branched; bracts sessile, ovate, narrow-ovate or narrow-lanceolate, 0.5-0.6 mm long, 0.2-0.35 mm broad, brown, glabrous, apex acute or attenuate. Pistillate flower 5-merous, not seen; persistent tepals 5, lanceolate-linear or narrowlanceolate, 0.7-1.5

收稿日期：2013-03-14 修回日期：2013-04-16

作者简介：王文采(1926-)男,山东掖县人,研究员,中国科学院院士,著名植物分类学家,长期从事有花植物分类学研究。




图1 柔毛赤车 A. 雌茎上部; B. 果序; C. 宿存花被片和瘦果; D. 瘦果。(来自模式标本)

Fig. 1 *Pellionia mollissima* A. Upper part of pistillate stem; B. Inflorescence; C. Persistent tepals and achene; D. Achene. (from holotype)

long 0.3–0.4 mm broad, brown, at the middle part dark-brown, glabrous, apex acute. Achenes bilaterally compressed, biconvex, broad-elliptic, ca. 1 mm long, 0.6 mm broad, longitudinally lineolate and very minutely tuberculate, brown, apex with white membranous

remnants of stigma ca. 0.2 mm long.

多年生草本, 当干燥时变为黑色。茎高约 65 cm, 近基部粗 6 mm, 上部密被短柔毛(毛长 0.2 mm), 下部无毛并具 10 条浅纵沟。叶无柄; 叶片薄纸质, 斜倒披针形或长椭圆形, 长 13~17 cm, 宽 2~

4.8 cm, 顶端长渐尖, 基部狭侧长渐狭, 宽侧钝或耳形, 边缘具小牙齿, 两面极密被短柔毛(毛长0.2~0.5 mm), 半离基三出脉, 侧脉在叶狭侧3条, 在宽侧5条, 钟乳体密集, 杆状, 长0.2~0.5 mm; 托叶膜质, 狹披针形或条形, 长9~11.5 mm, 宽2~2.8 mm, 背面被短柔毛。雄聚伞花序未见。雌聚伞花序单生叶腋, 小, 长和宽均为3 mm, 无梗, 下部自基部起有4条很短分枝; 苞片无柄, 卵形、狭卵形或狭披针形, 长0.5~0.6 mm, 宽0.2~0.35 mm, 褐色, 无毛, 顶端急尖或渐狭。雌花五基数, 未见; 宿存花被片5, 披针状条形或狭披针形, 长0.7~1.5 mm, 宽0.3~0.4 mm, 褐色, 中央暗褐色, 无毛, 顶端急尖。瘦果两侧扁, 双凸, 宽卵圆形, 长约1 mm, 宽0.6 mm, 具纵列短线纹和极小瘤状突起, 褐色, 顶端具长约0.2 mm的柱头膜质残遗部分。

广西(Guangxi): 龙州(Longzhou), 金龙(Jinlong), alt. 250 m, 石灰岩山林中(under forest in limestone hill), 2003-09-15, 稅玉民, 陈文红(Y. M.

Shui & W. H. Chen) B2003-038(holotype, PE)。

本种的突出特征是瘦果具纵列短线纹和小瘤状突起, 据此特征即可与赤车属的其他所有种相区别; 在此属其他种, 瘦果通常具瘤状突起, 在少数种光滑或具不规则条纹(marmorate)或短线纹(lineolate)(Schröter, 1936; 王文采, 1995)。此外, 本种的另一特征是叶两面均密被短柔毛, 据此一特征即可与我国赤车属的其他所有种相区别; 在我国此属其他种, 叶两面均有多少稀疏的毛, 或只上面或只下面有毛, 或两面均无毛(王文采, 1995)。

致谢 孙英宝先生为本文绘图, 深表感谢。

## 参考文献:

- 王文采. 1995. *Pellionia* [M] // 中国植物志. 北京: 科学出版社, 23(2): 160-187
- Schröter H, Winkler H. 1936. *Elatostema* subgen. [M] // *Pellionia*. Monographie der Gattung *Elatostema* s. l. Repert. Sp. Nov. Regn. Veg. Beih, 83(2): 2-92
- Hulme PE. 2008. Phenotypic plasticity and plant invasions: is it all Jack? [J]. *Funct Ecol*, 22(1): 3-7
- Lichtenthaler HK, Burkart S. 1999. Photosynthesis and high light stress [J]. *Bulg J Plant Physiol*, 25(3-4): 3-16
- Lowe S, Browne M, Boudjelas S, et al. 2000. 100 of the World's Worst Invasive Alien Species [M] // A selection from the Global Invasive Species Database. Auckland: New Zealand.
- Prentis PJ, Wilson JRU, Dormontt EE, et al. 2008. Adaptive evolution in invasive species [J]. *Trends Plant Sci*, 13(6): 288-294
- Sakai AK, Allendorf FW, Holt JS, et al. 2001. The population biology of invasive species [J]. *Ann Rev Ecol Syst*, 32: 305-332
- Sloop CM, Ayres DR, Strong DR. 2009. The rapid evolution of self-fertility in *Spartina* hybrids (*Spartina alterniflora* × *foliosa*) invading San Francisco Bay, CA [J]. *Biol Inv*, 11(5): 1131-1144
- Urbanska KM, Hurka H, Landolt E, et al. 1997. Hybridization and evolution in *Cardamine* (Brassicaceae) at Urnerboden, central Switzerland: Biosystematic and molecular evidence [J]. *Plant Syst Evol*, 204(3-4): 233-256
- Wang CD(王程栋), Wang SS(王树声), Shen GM(申国明), et al. 2007. A study of cell ultrastructure of tobacco leaf with different maturity grades(不同成熟度烟草叶片细胞超微结构变化规律研究) [J]. *Chin Tobac Sci*(中国烟草科学), 1(3): 30-34
- Wu W, Zhou RC, Ni GY, et al. 2013. Is a new invasive herb emerging? Molecular confirmation and preliminary evaluation of natural hybridization between the invasive *Sphagneticola trilobata* (Asteraceae) and its native congener *S. calendulacea* in South China [J]. *Biol Inv*, 15(1): 75-88
- Xia MY(夏明月), Chai Y(柴岩), Feng BL(冯佰利), et al. 2009. Ultrastructure changes of different leaf positions during different maturity grades in mung bean(绿豆开花后不同节位叶片衰老的超微结构研究) [J]. *Acta Agric Bor-Osc Sin*(西北农业学报), 31(3): 131-135
- Yang XH(杨兴洪), Chen CR(陈翠容), Shi P(施培). 2000. Effects of shading on the anatomical structure of cotton leaf and stem(遮荫对棉花茎叶解剖结构的影响) [J]. *J Shandong Agric Univ: Nat Sci Ed*(山东农业大学学报·自然科学版), 31(4): 373-377
- Yu LF(于凤龙), An FQ(安福全), Li FH(李富恒), et al. 2011. Effect of low light and low temperature on anatomical structure of summer squash leaves(低温弱光对西葫芦叶片解剖结构的影响) [J]. *J Changjiang Veg*(长江蔬菜), 6(6): 44-46
- Zhang SR(张守仁), Ma KP(马克平), Chen LZ(陈灵芝). 2002. Tempo-spatial variations in stomatal conductance, aperture and density of *Ligustrum sinense* acclimated to different light environments(适应不同光环境小蜡叶片气孔导度、气孔开度和气孔密度的时空变化) [J]. *Acta Bot Sin*(植物学报), 44(10): 1225
- Zhang ZX(张振贤), Guo YK(郭延奎), Zou Q(邹琦). 1999. Effects of shading on ultrastructure of chloroplast and microstructure of ginger leaves(遮荫对生姜叶片显微结构及叶绿体超微结构的影响) [J]. *Acta Hortic Sin*(园艺学报), 28(2): 28-32